

Animal welfare, etológia és tartástechnológia



Animal welfare, ethology and housing systems

Issue 3

Volume 9

Különszám/Special Issue

Gödöllő
2013



VARIANCES OF DIRECT AND MATERNAL GENETIC EFFECTS FOR MILK YIELD AND IT COMPOSITION IN A HERD OF FRIESIAN COWS IN EGYPT

Mostafa¹, M.A., Beder², E.A., Khattab³, A.S., Tözser⁴, J

¹Animal production Department, Faculty of Agriculture, Mansour University,

²Animal Production Research Institute, Ministry of Agriculture, Dokki, Cairo, Egypt,

³Animal production Department, Faculty of Agriculture, Tanta Univ., Egypt,

⁴Institute of Animal Husbandry, Szent Istvan University, H- 2013 Gödöllő Páter K. u. 1., Hungry
Adelkhattab@yahoo.com

Abstract

A total of 3748 normal lactation records of Friesian cows kept at Sakha farm, belonging to Animal production Research Institute, Ministry of Agriculture, Dokki, Cairo, Egypt during the period from 2000 to 2006 were used in the present study. Data was analyzed by using Multiple Traits Derivative Free Restricted Maximum Likelihood (MTDFREML) according to Boldman et al. (1995) using repeatability animal model. Two models were used. Model 1 includes month and year of calving and parity as fixed effects and direct genetic, maternal genetic, covariance between direct and maternal genetic and residual as random effects. Model 2 is similar to Model1, while additive maternal and covariance between direct and maternal effects were omitted from the model.

Estimates of heritability for 305 day milk yield, lactation period, fat % and protein % were 0.33, 0.02, 0.16 and 0.64, respectively for model 1, while the values were 0.34, 0.03, 0.17 and 0.65, respectively as estimated from model 2. The removal of additive maternal genetic effects and covariance between direct and maternal genetic effects from the model increased estimates for heritability of additive genetic effects by 0.01, for the four traits. Therefore, the additive maternal genetic effect and covariance between direct and maternal genetic effects do not seem to make important contributions to the phenotypic variance for milk yield and it is composition.

Introduction

Maternal effects have been defined as any influence from a dam on it is offspring, excluding the effects of directly transmitted genes that affect performance of the offspring (*Legates*, 1972). In mammals, environmental variation in the offspring is partially due to genetic variation of some other traits from the dams, such that quantitative traits can be influenced by two genetic components, animal genotype (direct genetic effect) and dam genotype (maternal genetic effects). The maternal genetic effects are not important for yield traits of dairy cattle (*Schutz* et al., 1992 ; *Albuquerque* et al., 1998 and *Khattab* et al., 2005). *Khattab* et al. (2005) with 2095 lactation records of Friesian cows in Egypt, found that means estimates of the effects of maternal genetic variances and direct maternal covariances as a fractions of phenotypic variances for 305 d MY were 0.01 and 0.02, respectively. The objectives of the present study were to quantify the contribution of additive direct and maternal genetic effects to phenotypic variance of milk yield and milk composition in a herd of Friesian cows in Egypt.



Material and methods

Source of data

Data used in the present study were collected from the history sheets of Friesian cows maintained at Sakha Farm, belonging to Animal production research Institute, Ministry of Agriculture, Dokki, Cairo, Egypt. The data comprised 3748 normal lactation records spread over the period from 2000 to 2006, representing in parities from 1 to 6. Abnormal records of cows affected by diseases such as mastitis and udder troubles or reproductive disorders were excluded. Animals were grazed on Egyptian clover (*Trifolium alexandrinum*), berseem, during December to May. They were fed concentrate mixture along with wheat or rice straw and limited amount of clover hay when available during the rest of the year. Cows producing more than 10 kg a day and cow a pregnant in the last two months were supplemented with extra concentrate. Heifers were first inseminated at 18 months of age. In subsequent lactations, cows were initially inseminated 60- 70 days postpartum. Cows were machined twice a day. Sires which had 5 daughters or more were included in the genetic analysis. Cows were artificially inseminated randomly to ensure that no specific sires were bred to certain good or poor dams. Traits studied are 305 day milk yield, lactation period, fat % and protein %. Fat and protein percentages estimated using system of Milko – Scan 130 series. Type 10900.

Analysis

The records were analyzed by Multiple Trait Derivative Free Restricted Maximum Likelihood (MTDFREML) according to *Boldman et al* (1995), using repeatability animal model. Two models were used, *Model 1*, includes month of calving, year of calving and parity as fixed effects and direct genetic, maternal genetic, covariance between direct and maternal genetic, permanent environmental and residual as random effects. *Model 2* is similar to *Model 1*, but excluding additive maternal and covariance between additive direct and maternal effects. *Table 1* shows the data structure considered in the analysis.

Table 1: Structure of data used in analysis, means, standard deviation (SD) and coefficient of variability (CV %) for 305 day milk yield (305 d MY), lactation period (LP), fat % (F %) and protein % (P %)

Traits	Mean	SD	CV %
305 d MY, kg	3558	119	40
LP, d	301	172	48
F %	3,7	0,82	22
P %	2,60	0,51	20



Observations

No. of records	3748
No. of sires	50
No. of cows	428
No. of dams	191
Animals in relationship matrix (A^{-1})	669
Mixed Model Equations (MME)	5460
No. of iterations	11789

Results and discussions

Means of 305 d MY, LP, F % and P % , standard deviations (SD) and coefficients of variability (CV %) are presented in table 1. Means of 305 d MY, LP, F % and P % were 3558 kg, 301 d, 3.7 % and 2.60 % (Table 1). The present means of 305 d MY and LP were higher than those found by *Yener et al.* (2006) working on another set of Friesian cows in Egypt, reported that means of 305 d MY and LP were 2806 kg and 293 d, respectively. While the present means were lower than those reported by *Khattab and Atil* (1999) working on another set of Friesian cows in Egypt, found that means of 305 d MY and LP were 3709 kg and 367 d, respectively. Overall mean of F % was similar to that reported by *Albuquerque et al.* (1998) (3.7) working on Holstein Friesian. While The present mean of F % was lower than those found by *Islam et al.* (2008) (4.28%). The overall mean of P % was lower than those reported by *Jairath et al.* (1995) (3.15) on Canadian Holstein and *Islam et al.* (2008) (3.11) on Holstein Friesian. The difference between the present estimates and those reported by other studied could be due to differences in climatic and management. Herds could be possibly the genetically different or be caused by different models used.

The large CV % value for 305 d MY (40 %) reflects a great variation between individuals in such an important productive trait.

Variance components and means for variance and covariance components as ratio of phenotypic variances from model 1 and 2 are in *Table 2*. The variance for additive maternal genetic effects were 4, 2.5 and 7 % of phenotypic variance for 305 d MY, LP, F % and p %, respectively and the covariance between maternal and direct genetic effects were 0.07, 0.01, 1 and 2 % for the same traits respectively. Thus, these effects did not contribute importantly to phenotypic variance. Similar results were obtained by *Albuquerque et al.* (1998) analysis milk traits (milk, fat yield and fat %) reported that the variance of additive maternal genetic effects varied from 0.8% to 1.0 % of phenotypic variance and the covariance between maternal and direct genetic effects varied from 0.7 % to 2.5 %. Also, *Schutz et al.* (1992) using an animal model estimated small values for ratios of variance for maternal genetic effects to phenotypic variance for milk yield (2.58 %) and fat % (6.5 %0 and for covariance between maternal and direct effects for milk yield (5.99 %). In addition, *Khattab et al.* (2005) with another set of Friesian cows in Egypt, found that the variance for additive maternal genetic effect and covariance between maternal and direct genetic effects for 305 d MY are 1 % and 2 %, respectively.



Maternal genetic effects have been described in domestic mammals such as swine (Southwood and Kennedy, 1990) and beef cattle (Dodenhoff et al, 1990) and represent an environmental effect on the growth of offspring from birth to weaning. However, dairy calves are separated from their dams at birth so that the influence of the dam would be only through intrauterine environmental (Albuquerque et al., 1998). In addition, uterine environmental and the maternal colostrums may also contribute to the maternal genetic effects (Khattab et al, 2005).

Table 2: Phenotypic and genetic variance and covariance for 305 day milk yield (305 d MY), lactation period (LP), fat percent (F %) and protein percent (P %) using two models

Model	g^2	m^2	gm	e^2	Total
Model 1					
305 d MY	25.52 (0.33)	2.91 (0.04)	-0.58 (0.007)	49.82 (0.64)	77.67
LP	2.58 (0.02)	1.80 (0.02)	-0.02 (0.001)	113.47 (0.96)	117.83
F%	4.71 (0.16)	1.41 (0.05)	-0.36 (0.01)	24.14 (0.81)	29.90
P%	9.40 (0.64)	0.98(0.07)	-0.26 (0.02)	4.47 (0.31)	14.59
Model 2					
305 d MY	26.40 (0.34)			50.20 (0.66)	76.60
LP	3.10 (0.03)			114.47(0.97)	117.57
F%	5.10 (0.17)			25.10 (0.83)	30.20
P%	11.20 (0.65)			5.90 (0.35)	17.10

G^2 = additive genetic effect, m^2 = additive maternal genetic effect, gm= covariance between direct and maternal.

Heritability estimates for 305 d MY, LP, F % and P % as estimated from model 1 (Full model) are 0.33 ± 0.02 , 0.02 ± 0.01 , 0.16 ± 0.04 and 0.64 ± 0.13 , respectively (Table 3), while the corresponding values as estimated from model 2 are 0.34, 0.03, 0.17 and 0.65, respectively (Table 2). The removal of additive genetic maternal effects and covariance between direct and maternal effects from the model (Model 2) increased estimates of heritability of direct genetic effects by 0.01 for all traits studied (Table 2). Similar results are reported by Albuquerque et al, (1998) found that heritability estimates of direct genetic effects increased by 0.014, 0.021 and 0.046 for milk yield, fat yield and fat %, respectively, when additive genetic maternal effects and covariance between direct and maternal genetic effects are removed from the analysis. The same authors also concluded that, some confounding between direct and maternal genetic effects should be expected because the dam that contributes the maternal genetic effect also transmits half of her genetic value for direct effects in her daughter. Some cows do not have lactating daughters, and many sires do not have any lactating granddaughters. Also, Schutz et al, (1992) and Khattab et al, (2005) arrived at the same conclusion, reported that maternal genetic effects are not important for yield traits of dairy cattle.



Table 3: Estimates of heritability (direct and maternal) and genetic correlation among 305 d MY, LP, F % and P % as estimated from model 1*.

	g^1	g^2	g^3	g^4	m^1	m^2	m^3	m^4
g^1	0.33±0.02							
g^2	0.16±0.01	0.02±0.01						
g^3	-0.10±0.01	-0.12±0.02	0.16±0.04					
g^4	-0.13 ±0.01	-0.18±0.02	0.45±0.14	0.64±0.13				
m^1	-0.07±0.02	-0.13±0.03	0.15±0.10	-0.52±0.18	0.04±0.01			
m^2	0.06±0.02	C	-0.19±0.18	-0.11±0.07	0.06±0.00	0.01±0.00		
m^3	-0.07 ±0.01	-0.09±0.02	-0.14±0.07	-0.16±0.04	-0.17±0.01	-0.24±0.01	0.05±0.01	
m^4	-0.18±0.02	-0.08±0.02	0.09±0.01	-0.07±0.03	0.37±0.01	0.42±0.01	-0.30±0.10	0.06±0.01

*

305 d MY, LP, F %, P%, $g^1, g^2, g^3, g^4, m^1, m^2, m^3$ and m^4 are 305 day milk yield, lactation period, fat %, protein %, direct heritability for 305 d MY, direct heritability for LP, direct heritability for F %, direct heritability for P %, maternal heritability for 305 d MY, maternal heritability for LP, maternal heritability for F % and maternal heritability for p %, respectively.

Direct heritability estimates for 305 d MY, LP, F % and P % as estimated from model 1 (full model) are 0.33, 0.02, 0.16 and 0.64, respectively (Table 3). The present estimates are within the range of recorded value with an animal model (i.e., *Albuquerque et al., 1998; El- Arian et al., 2003 ; Yener et al., 2006 and Usman et al., 2012*) and ranged from 0.07 to 0.33. The moderate estimates of heritability for 305 d MY and F % and high estimate of heritability for P %, suggests that more efforts could be made to bring about improvement milk yield and it is composition through individual selection as well as better managerial practices. Low heritability estimates for LP indicate that this trait is affected mainly by environmental factors.

The results in table 3 show that the genetic correlation between 305 d MY and LP was positive and being 0.16±0.01 and in the desirable direction indicating that high yielding cows are also have the longer LP. Whereas the genetic correlation between 305 d MY and each of F % and P% were negative and being -0.10 ± 0.01 and -0.13 ± 0.01 , respectively. Also, the genetic correlation between LP and each of F and P % were negative and being -0.12 ± 0.02 and -0.18 ± 0.02 , respectively (Table 3). The present results indicated that selection for milk yield and lactation period would lead to slight decrease fat and protein percent. While, genetic correlation between F and P % was positive (0.45±0.14). Similar results are reported by *El- Arian et al. (2002), Yener et al. (2006) and Usman et al. (2012)*. Also, maternal genetic correlation among milk traits studied are in most cases negative and small.

According to small amount of additive maternal genetic effects for 305 d MY, LP, F % and P % (Table 2), it could be concluded that the additive maternal genetic effects and the covariance between additive maternal and direct genetic effects do not seem to make important contributions to the phenotypic variance of milk traits, probably because the important environmental influence of the dams on their calves is from conception to birth.



References

- Albuquerque, L. D.; Keown, J.F.; Van Vleck, L.D* (1998) Variances of direct genetic effects, maternal genetic effects and cytoplasm inheritance effects for milk yield, fat yield and Fat percentage. *J. Dairy Sci.*, Vol 81: 544- 549.
- Boldman, K.G.; Kriese, L. A.; Van Vleck, L.D.; Kachman, S. D.*(1995) A manual for use of MTDFREML , Department of Agriculture/ Agricultural Research Service, Lincoln, 120 pp.
- Dodenhoff, J.; Van Vleck, L.D.; Gregory, K.E.* (1999) Estimation of direct, maternal and maternal genetic effects for weaning weight in several breeds of beef cattle. *J. Anim.*
- El- Arian, M.N.; El- Awady, H.M.; Khattab, A.S.* (2003) Genetic parameters and breeding values of milk production traits by using animal model in a herd of Holstein Friesian Cows in Egypt. 53rd Annual Meeting of the European Association for Animal Production, Cairo, Egypt, 1-4 September, 2003.
- Islam, M.R.; Hassan, M.N.; Siddiki, M.S.R.; Khan, M.A.S.; Islam, M.A.* (2008) Determination of quality of milk from different genotype of dairy cows. Bangladesh. *J. Anim. Sci.*, 37:53.
- Jairath, L.K.; Hayes, J.F.; Cue, R.I.*(1995) Correlations between first lactation and lifetime performance traits of Canadian Holsteins. *J. Dairy Sci.*, 78:438.
- Khattab, A.S.; Atil, H.*(1999) Genetic study of fertility traits and productive in a local Friesian cattle in Egypt. *Pakistan J. of Biol.* 2:1178.
- Khattab, A.S.; Atil, H.; Badawy L.* (2005) Variances of direct and maternal genetic effects for milk yield and age at first calving in a herd of Friesian cattle in Egypt. *Arch. Tierz. Dummerstorf.* 48:24-31.
- Legates, J.E.* (1972) The role of maternal effects in animal breeding. IV. Maternal effects in laboratory species. *J. Anim. Sci.*, 35: 1294-1302.
- Schutz, M. M.; Freeman, A.E.; Beitz, D.C.; Mayfield, J.E.* (1992) The importance of maternal lineage on milk yield traits of dairy cattle. *J. Dairy Sci.*, 75:1331-1341.
- Southwood, O. I.; Kennedy, B.W.* (1999) Estimation of direct and maternal genetic variance for litter size in Canadian Yorkshire and Landrace Swine using an animal model. *J. Anim. Sci.*, 68:1841-1847.
- Usman, T.; Guo, G.; Suhail, S.M.; Wang, Y.* (2012) Performance traits study of Holstein Friesian cattle under subtropical conditions. *J. of Animal and Plant Sci.*, 22:92.
- Yener, S.M.; Akman, N.; Khattab, A.S.; Hussein, A.* (2006) Genetic aspects for productive reproductive traits for Friesian cows in Egypt. 57th Annual meeting of the European Association for Animal Production, Antalya, Turkey.